

NOTE XIX.

TWO NEW SPECIES OF CICINDELIDAE

BY

Dr. WALTHER HORN.

Tetracha lateralis, n. sp.

Species intermedia inter *T. spinosam* Brll. et *T. Steinheili* m.; differt ab illa oculis minoribus magisque convexis; prothorace et antice et praecipue postice constricto, ut latera non recta videantur; impressionibus longitudinalibus elytrorum paullo minus distinctis, angulo suturali acuto (non spinoso!), margine epipleurisque similimodo sed profundius sinuato-curvatis, macula flava apicali multo breviore, 3 primis sternitis abdominalibus lateraliter metallicis (non testaceis), ceteris discoque (hoc levissime metallice induito) infuscatis; tarsis (posticis) brevioribus. Corpore supra obscure. — Long. 13 mm.

1 ♀; Amazon. — Coll. mea.

Differt a *T. Steinheili* m. oculis majoribus minusque convexis; prothorace ad angulos anticos minus dilatato, postice minus constricto; elytris in medio et pone humeros magis dilatatis, ad apicem magis applanatis, apice singulo non rotundato, sculptura omnino differente (ante medium minus grosse punctatis, postea sat dense granulatis — non rare punctatis — interstitiis non glabro-politis etc.); tarsis (posticis) longioribus et angustioribus; corpore supra non cyanescente; macula flava apicali elytrorum multo angustiore (antice angustata!), 2°—4° antennarum articulis solummodo intus nigro-annulatis; femoribus distaliter non macula infuscata ornatis etc.

T. spinosa Brll. was formerly a quite isolated species in the whole genus, now it is connected with *T. cibrata* Steinh. etc. by *T. lateralis* m. and *T. Steinheili* m.

Odontochila curvipenis, n. sp.

Od. margineguttatae Dej. simillima, differt labri parte media latius et paullo longius producta, dente centrali nullo (hoc loco leviter exciso), ad medianum basim infuscato (non metallico); prothoracis lateribus paullo minus rotundatis, pronoto grossius sculpto (transversaliter rugatulo); elytris praecipue ante medium paullo grossius punctatis, penis apice curvato-acuminato (nec recto nec incrassato); penultimo palporum maxillarium articulo, trochanteribus, primo antennarum articulo testaceis (2° — 4° cyaneis saepe ex parte testaceo-rufescensibus); femoribus et tibiis testaceis saepe leviter variis modis rufo-violaceo-indutis, tarsis cyaneis, corpore subtus cyanescente (abdomine hinc inde virescente). — Long. 9— $9\frac{1}{2}$ mm. (sine labro).

♂♂; »Amazon» et »Maices». — Coll. mea.

This plain-looking species is very remarkable by the form of the tip of the penis. The sculpture, coloration etc. show also some characteristical features.

Berlin, May 1905.